

## Apoptotic and cytostatic actions of maslinic acid in colorectal cancer cell lines through possible IKK- $\beta$ inhibition


Kai Xin Ooi<sup>1</sup>, Menaga Subramaniam<sup>1</sup>, Lai Yen Fong<sup>2</sup>, Hoe Han Goh<sup>3</sup>, Soo Beng Alan Khoo<sup>4</sup>, Yang Mooi Lim<sup>1,2\*</sup>

<sup>1</sup>Centre for Cancer Research, Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, LOT PT 21144. Jalan Sungai Long, Bandar Sungai Long, Cheras 43000 Kajang, Selangor, Malaysia

<sup>2</sup>Department of Pre-Clinical Sciences, Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, LOT PT 21144. Jalan Sungai Long, Bandar Sungai Long, Cheras 43000 Kajang, Selangor, Malaysia

<sup>3</sup>Institute of Systems Biology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor Darul Ehsan, Malaysia

<sup>4</sup>Molecular Pathology Unit, Cancer Research Centre, Institute for Medical Research, Kuala Lumpur, Malaysia



**Supplementary Figure 1** Chemical structure of maslinic acid [(2 $\alpha$ ,3 $\beta$ )-2,3-dihydroxyolean-12en-28-oic acid].